

Appendix

Appendix Table 1: Atrial fibrillation (AF) and oral anticoagulant use (OAC) by district.

DHB of domicile	No AF	Known AF on OAC	Known AF not on OAC	AF Dx post-stroke	Unknown	Total	% AF	% known AF not on OAC/all stroke	% known AF not on OAC/known AF
Northland	58	12	15	12	0	97	40%	15%	56%
Waitematā	157	27	22	14	0	220	29%	10%	45%
Auckland	77	21	5	2	0	105	27%	5%	19%
Counties Manukau	117	53	8	10	1	189	38%	4%	13%
Waikato	117	42	13	8	9	189	38%	7%	24%
Lakes	62	16	11	9	5	103	40%	11%	41%
Bay of Plenty	108	37	20	13	0	178	39%	11%	35%
Tairāwhiti	18	12	0	3	2	35	49%	0%	0%
Taranaki	57	17	18	4	1	97	41%	19%	51%
Hawke's Bay	57	33	3	2	0	95	40%	3%	8%
MidCentral	67	16	7	5	8	103	35%	7%	50%
Whanganui	50	7	7	4	2	70	29%	10%	30%
Capital and Coast	93	17	13	18	1	142	35%	9%	43%
Hutt	71	7	11	3	0	92	23%	12%	52%
Wairarapa	27	14	1	0	1	43	37%	2%	7%
Nelson Marlborough	35	10	7	0	0	52	33%	13%	41%
West Coast	15	5	4	1	2	27	44%	15%	44%

Appendix Table 1 (continued): Atrial fibrillation (AF) and oral anticoagulant use (OAC) by district.

Canterbury	162	52	39	22	3	278	42%	14%	43%
South Canterbury	25	11	3	2	0	41	39%	7%	21%
Southern	124	33	16	8	1	182	32%	9%	33%
Overseas	12	0	1	1	0	14	14%	7%	50%
Total	1,509	442	224	141	36	2,352	36%	10%	27%