




Appendix 1: Age- and sex-standardised incidence rate (SIR) and absolute numbers of annual cases for the top 10 most commonly diagnosed cancers for Pacific **females** between 2007–2019 (top), along with the age- and sex-standardised rate difference (SRD) between Pacific females and European females (bottom).




Appendix 2: Age- and sex-standardised incidence rate (SIR) and absolute numbers of annual cases for the top 10 most commonly diagnosed cancers for Pacific males between 2007–2019 (top), along with the age- and sex-standardised rate difference (SRD) between Pacific males and European males (bottom).



Appendix 3: Age- and sex-standardised mortality rate (SMR) and absolute numbers of cases for the top 10 most common causes of cancer death for Pacific **females** between 2007–2018 (top), along with the age- and sex-standardised rate difference (SRD) between Pacific females and European females (bottom).



Appendix 4: Age- and sex-standardised mortality rate (SMR) and absolute numbers of cases for the top 10 most common causes of cancer death for Pacific males between 2007–2018 (top), along with the age- and sex-standardised rate difference (SRD) between Pacific males and European males (bottom).



Appendix 5: Annual cancer cases, incidence rates and mortality rates for Pacific and European peoples. SIR: Standardised incidence rate; SMR: standardised mortality rate.

Cancer name	Incidence				Mortality			
	Cases/year		SIR (95% CI)		Deaths/year		SMR (95% CI)	
	Pacific	European	Pacific	European	Pacific	European	Pacific	European
Bladder	8	353	3.9 (1.2–6.6)	5.4 (4.7–6)	4	189	-	-
Bone and cartilage	5	25	1.5 (0.1–2.9)	0.9 (0.5–1.3)	2	13	-	-
Breast	166	2,370	126.9 (107.3–146.5)	114.9 (109.5–120.3)	34	507	26.5 (17.5–35.5)	19.4 (17.3–21.4)
Cervix	14	97	10.3 (4.8–15.7)	7 (5.4–8.5)	6	34	4.3 (0.7–7.8)	1.5 (0.9–2.1)
Colorectal	73	2,677	32 (24.5–39.5)	45.9 (43.9–48)	29	1,100	12.9 (8.1–17.7)	16.6 (15.5–17.8)
Eye, brain and CNS	18	311	6.4 (3.4–9.4)	7.9 (6.9–8.9)	9	235	3.3 (1–5.5)	5.4 (4.6–6.2)
Gallbladder and biliary tract	12	101	5.4 (2.2–8.5)	1.6 (1.2–2)	6	69	2.9 (0.6–5.2)	1 (0.7–1.3)
Head and neck	29	442	12.2 (7.6–16.7)	9.6 (8.6–10.6)	10	138	4.4 (1.6–7.1)	2.4 (1.9–2.8)
Hodgkin's lymphoma	7	78	2.3 (0.5–4.1)	2.9 (2.2–3.6)	1	16	-	-
Ill-defined, secondary or unspecified	20	356	9 (5–13)	5 (4.4–5.6)	15	328	6.8 (3.3–10.3)	4.3 (3.8–4.9)
Kidney	20	440	8.2 (4.5–11.9)	9.3 (8.3–10.3)	4	161	-	-
Leukaemia	33	534	13.1 (8.5–17.7)	11 (9.9–12.1)	15	261	6.2 (2.9–9.4)	4.1 (3.5–4.7)
Liver	31	190	14.3 (9.2–19.4)	3.4 (2.9–4)	21	157	9.9 (5.6–14.2)	2.6 (2.2–3.1)
Lung	104	1,520	49.1 (39.6–58.7)	24.6 (23.2–26)	72	1,229	34.2 (26.2–42.3)	19.2 (18–20.4)
Melanoma	8	2,493	3.3 (0.9–5.6)	56.2 (53.6–58.8)	2	328	-	-
Mesothelioma and soft tissue	14	207	5.8 (2.7–8.9)	4.3 (3.6–5)	6	142	2.5 (0.4–4.6)	2.5 (2–3)
Myeloma	22	279	10 (5.7–14.2)	4.9 (4.2–5.5)	10	147	4.5 (1.6–7.4)	2.1 (1.7–2.5)

Appendix 5 (continued): Annual cancer cases, incidence rates and mortality rates for Pacific and European peoples. SIR: Standardised incidence rate; SMR: standardised mortality rate.

Cancer name	Incidence				Mortality			
	Cases/year		SIR (95% CI)		Deaths/year		SMR (95% CI)	
	Pacific	European	Pacific	European	Pacific	European	Pacific	European
Non-Hodgkin's lymphoma	32	682	13.5 (8.8–18.3)	13.7 (12.5–14.9)	11	254	4.6 (1.7–7.4)	3.9 (3.4–4.5)
Oesophagus	8	250	3.7 (1.1–6.4)	4.1 (3.5–4.7)	5	210	2.4 (0.3–4.6)	3.3 (2.8–3.8)
Ovary	24	255	18.4 (11–25.8)	10 (8.6–11.5)	10	186	7.7 (2.8–12.5)	6 (5–7.1)
Pancreas	22	433	9.7 (5.6–13.8)	7 (6.3–7.8)	15	385	6.8 (3.3–10.3)	6 (5.3–6.7)
Prostate	105	2,899	108.7 (87.8–129.7)	110.4 (106.1–114.7)	18	555	19.8 (10.6–29)	14.6 (13.3–15.9)
Small intestine	6	79	2.7 (0.5–4.8)	1.5 (1.1–1.9)	2	35	-	-
Stomach	37	244	16.3 (11–21.7)	4.3 (3.7–4.9)	23	191	10.2 (5.9–14.4)	3.1 (2.6–3.6)
Testis	8	107	4.9 (1.4–8.4)	9.9 (8–11.8)	0	5	-	-
Thyroid and endocrine	31	175	11.6 (7.4–15.7)	5.5 (4.6–6.4)	3	24	-	-
Uterus	79	333	61 (47.3–74.6)	13.1 (11.5–14.7)	16	79	12.9 (6.4–19.3)	2.3 (1.7–2.9)