Figure 1: Community systemic antibiotic dispensing between 2010 and 2019 by defined daily doses per 1,000 inhabitants per day (DIDs) (a), and antibiotic courses dispensed per 1,000 population per year (b).



Figure 2: Percentage of community systemic antibiotic dispensing by the World Health Organization AWaRe groups in Aotearoa New Zealand during 2010-2019.





Figure 3: The number of systemic antibiotic courses dispensed per 1,000 inhabitants per year in the Aotearoa New Zealand community between 2010 and 2019 by age group.



Figure 4: Age-standardised rates of community systemic antibiotic dispensing between 2010 and 2019 by ethnicity measured by defined daily doses per 1,000 inhabitants per day (DIDs) (a), and antibiotic courses dispensed per 1,000 population per year (b).



Figure 5: Age-standardised rates of community systemic antibiotic dispensing between 2010 and 2019 by New Zealand Deprivation (NZDep) quintiles (quintile 1 = least deprived, quintile 5 = most deprived). Measured by the number of defined daily doses per 1000 population per day (DIDs) (a), and the number of antibiotic courses dispensed per 1,000 inhabitants per year (b).



(a) y axis range is 16–24 DIDs and (b) y axis range is 600–1200 courses dispensed per 1,000 inhabitants per year.